

## NOTE

Masaru Miyata · Kazutaka Itoh · Sanro Tachibana

**Extractives of *Juniperus chinensis* L. I: Isolation of podophyllotoxin and yatein from the leaves of *J. chinensis***

Received: November 25, 1997 / Accepted: May 6, 1998

**Abstract** Two compounds, yatein and podophyllotoxin, were isolated from the chloroform-soluble fraction in the methanolic extractives of byakushin (*Juniperus chinensis* L.) leaves for the first time.

**Key words** *Juniperus chinensis* · Extractives · Lignans · Podophyllotoxin · Yatein

**Introduction**

This study is a continuation of our work on the production of biologically active substances by tissue cultures of trees<sup>1,2</sup> and the microbial and enzymatic conversion of extractives from trees.<sup>3–7</sup> We investigated the extractives of byakushin (*Juniperus chinensis* L.) in search of potential compounds for biotechnological production of biologically active substances.

*Juniperus chinensis* is an evergreen tree with scaly, partly needle, year-round foliage. It is grown in Japan, Korea, and China.<sup>8</sup> To date, only two studies have examined extractives from the leaves of this plant. One study, conducted by Sawada,<sup>9</sup> isolated two bisflavonoids (hinokiflavone and kayaflavone). The other study, by Fang et al.,<sup>10</sup> isolated a coumarin (umbelliferone), 13 lignans, and 2-arylpropane-1,3-diol. Fang et al. did not isolate podophyllotoxin or yatein from the leaves of kaizukaibuki (*J. chinensis* var. Kaizuka Hort.), a variety of byakushin.

Podophyllotoxin, a lignan with strong antileukemic and tumor-inhibiting effects, has been isolated from the leaves of *Juniperus sabina*, along with several podophyllotoxin-related compounds.<sup>11</sup> It is believed that podophyllotoxin

and related compounds are contained in *J. chinensis* leaves. Podophyllotoxin is a pharmaceutically valuable compound that has become commercially important as raw material for the semisynthesis of the antitumor agents etoposide and teniposide. Natural sources of podophyllotoxin are mainly species of *Podophyllum*, but a search for a new source of podophyllotoxin other than species of *Podophyllum* is required because large amounts of podophyllotoxin are needed as raw material for production of the antitumor agents. Isolation of podophyllotoxin and related compounds from the leaves of *J. chinensis* has not yet been reported. We describe herein the isolation of podophyllotoxin and yatein, a compound related to podophyllotoxin, from the leaves of *J. chinensis*.

**Results and discussion**

In the present study of *J. chinensis* leaves, two compounds were isolated: yatein (compound 1) and podophyllotoxin (compound 2). This is the first time that compounds (1) and (2) have been found in these sources. Their chemical structures are shown in Fig. 1. The structures were determined by ultraviolet (UV), proton nuclear magnetic resonance (<sup>1</sup>H-NMR), and carbon-13 nuclear magnetic resonance (<sup>13</sup>C-NMR) spectroscopy and mass spectrometry (MS).

Compound (1) was isolated as a colorless oil, C<sub>22</sub>H<sub>24</sub>O<sub>7</sub>, [ $\alpha$ ]<sub>D</sub><sup>25</sup> -25.0° (CHCl<sub>3</sub>), whose molecular ion peak (M<sup>+</sup>) was observed at m/z 400 in the mass spectrum. The UV spectrum of compound (1) had bands at  $\lambda_{\text{max}}^{\text{CHCl}_3}$ : 287 and 242 nm and showed a characteristic UV absorption spectrum of dibenzylbutyrolactone lignans.<sup>12</sup> The mass spectrum of the compound agreed well with the proposed structure.<sup>13</sup> No acetate was formed on acetylation of compound (1) with acetic anhydride and pyridine, which showed that compound (1) had no free phenolic or primary and secondary alcoholic hydroxyl groups. In the <sup>1</sup>H-NMR spectrum of the compound, signals of a 1,3,4-tri-substituted aromatic ring at 6.46, 6.47, and 6.69 ppm, a 1,3,4,5-tetra-substituted aromatic ring at 6.36 ppm, and two protons of a methylenedioxy at

M. Miyata · K. Itoh · S. Tachibana  
College of Agriculture, Ehime University, Matsuyama 790-8566,  
Japan

This report was presented at the 47th annual meeting of the Japan Wood Research Society, Kochi, 1997

5.93 and 5.94 ppm were observed. Furthermore, signals of four benzyl protons attached at the C-7' and C-7 positions at 2.48–2.63 ppm and 2.89–2.92 ppm, respectively, two protons of a methylene attached to the lactone oxygen at 4.18 and 3.88 ppm, two *trans*-protons attached at the C-8 and C-8' positions at 2.48–2.63 ppm, and three methoxyl groups at 3.82 ppm were also observed in the spectrum. MS and  $^1\text{H}$ -NMR spectra of compound (1) were in good agreement with those of authentic samples of yatein isolated from *Podophyllum hexandrum*.<sup>14</sup> The  $^{13}\text{C}$ -NMR spectrum well explained the structure of yatein.<sup>15</sup> The  $^{13}\text{C}$ -NMR assignments are shown in Table 1. Furthermore, the specific rotation ( $[\alpha]_{\text{D}}$ ) of compound (1) agreed with that of yatein isolated from *Libocedrus yateensis*.<sup>16</sup> Therefore, compound (1) was identified as yatein and was obtained in a yield of 0.0008% of the fresh leaves.

Compound (2) was isolated as white crystals,  $C_{22}H_{22}O_8$ , melting point (mp)  $181^{\circ}$ – $183^{\circ}C$ ,  $[\alpha]^{25}_D$   $-121.3^{\circ}$  ( $CHCl_3$ ), whose  $M^{+}$  was observed at  $m/z$  414 in the mass spectrum. The UV spectrum of compound (2) had bands at  $\lambda^{EtOH}_{max}$ : 291 and 219nm. The mass spectrum of compound (2) agreed well with the proposed structure.<sup>17</sup> In the  $^1H$ -NMR spectrum of the compound, signals of a 1,2,4,5-tetra-substituted aromatic ring at 6.51 and 7.11 ppm, a 1,3,4,5-tetra-substituted aromatic ring at 6.37 ppm, two protons of a methylenedioxy at 5.97 and 5.99 ppm, two methoxyl groups at 3.76 ppm, and a methoxyl group at 3.81 ppm attributed to the substituents at the C-3 and C-5 positions and the C-4 position, respectively, were observed. Furthermore, signals of two benzyl protons attached at the C-7 and C-7' positions at 4.62 and 4.77 ppm, two protons of a methylene attached to the lactone oxygen at 4.08 and 4.59 ppm, two *trans*-protons attached at the C-8 and C-8' positions at 2.7–2.9 ppm, and a proton of a hydroxyl group attached at the C-7' position at 2.17 ppm were also observed in the spectrum. MS and  $^1H$ -NMR spectra of compound (2) were in good agreement with those of authentic samples of podophyllotoxin purchased from Sigma Chemical Company.<sup>17</sup> The  $^{13}C$ -NMR spectrum well explained the structure of the compound.<sup>15</sup> The  $^{13}C$ -NMR assignments for compound (2)

Carbon <sup>b</sup>	Yatein	Podophyllotoxin
C-1	133.3	135.4
C-2	106.3	108.5
C-3	153.2	152.6
C-4	136.8	137.3
C-5	153.2	152.6
C-6	106.2	108.5
C-7	38.3	44.1
C-8	46.4	45.3
C-9	178.5	174.4
C-10	56.1	56.3
C-11	60.8	60.8
C-12	56.1	56.3
C-1'	131.5	133.1
C-2'	108.7	106.3
C-3'	147.9	147.7
C-4'	146.4	147.8
C-5'	108.3	109.8
C-6'	121.5	131.2
C-7'	35.2	72.8
C-8'	41.0	40.8
C-9'	72.0	71.3
C-10'	101.1	101.5

are shown in Table 1. The specific rotation ( $[\alpha]_D$ ) of compound (2) agreed with that of podophyllotoxin,<sup>18</sup> which was isolated from the podophyllin N.F. prepared from *Podophyllum peltatum*.<sup>19</sup>

On acetylation of compound (2) with acetic anhydride and pyridine, crystalline monoacetate (3), mp 207°–209°C, was obtained. The <sup>1</sup>H-NMR and MS spectra of compound (3) were in good agreement with those of authentic samples prepared from podophyllotoxin. The <sup>13</sup>C-NMR spectrum of compound (3) coincided with that of authentic podophyllotoxin monoacetate.<sup>15</sup> The mixed-melting-point test of compound (3) with authentic podophyllotoxin monoacetate proved the compound (3) to be identical with podophyllotoxin monoacetate. Therefore, compound (2) was identified as podophyllotoxin and was obtained in a yield of 0.0009% of the fresh leaves.

Podophyllotoxin and its related compound yatein were isolated for the first time from the leaves of *J. chinensis*. Podophyllotoxin has strong antileukemic and tumor-inhibiting activities.<sup>10</sup> Several studies have indicated that podophyllotoxin is contained in the leaves of *Juniperus sabina* (genus *Juniperus*, Cupressaceae)<sup>11</sup> and in the rhizomes of *Podophyllum* species.<sup>17,21</sup> The amounts of podophyllotoxin in *J. sabina* and *J. chinensis* are small, but species of *Podophyllum* produce large amounts of this lignan in their roots. Podophyllotoxin content is considered to be increased by tissue cultures of *J. chinensis*. We found that the content of podophyllotoxin could be increased by about 15 times that of the intact plant by callus cultures of *J. chinensis*.<sup>22</sup> Production of podophyllotoxin by callus cultures and cell suspension cultures of *J. chinensis* are now being conducted.

## Experimental

### Plant material

Fresh leaves (mixture of scaly and partly needle foliage) of *J. chinensis* were collected in May 1996 in Uwajima City, Ehime Prefecture.

### Extraction from *J. chinensis* leaves

Extraction from fresh leaves of *J. chinensis* (17 kg) was carried out twice for 4 days with methanol at room temperature. The methanolic extractives (1.09 kg) were suspended with water and successively extracted with *n*-hexane, chloroform, ethyl acetate, and *n*-butanol. The chloroform-soluble fraction gave a positive color reaction of *Podophyllum* lignans<sup>20</sup> with acetic acid and concentrated nitric acid (10:3 v/v) on a thin-layer chromatography (TLC) plate.

### Isolation of compounds (1) and (2) from chloroform-soluble fraction

The chloroform-soluble fraction (64.8 g) was separated into two fractions (fractions 1 and 2) by silica gel column chromatography with a chloroform-methanol stepwise elution. Fractions 1 and 2 were eluted out, in order. Fraction 1 was chromatographed on a silica gel column using a benzene-ethyl acetate stepwise elution and yielded fraction 1-1, containing a lignan (291 mg). Fraction 1-1 was rechromatographed on a preparative thin-layer chromatography (PTLC) with chloroform-acetone (65:35 v/v) and yielded compound (1) (133 mg).

Fraction 2 was chromatographed on a silica gel column using chloroform-ethyl acetate stepwise elution followed by benzene-ethyl acetate stepwise elution to afford fraction 2-1, containing a lignan (192 mg). Fraction 2-1 was rechromatographed on PTLC with chloroform-acetone (65:35 v/v) to give crude compound (2) (56.6 mg).

### Yatein

Compound (1) (yatein) was obtained as a colorless oil. *Analysis*. Calculated:  $C_{22}H_{24}O_7$ : C, 65.97; H, 6.04. Found: C, 65.82; H, 6.14.  $[\alpha]_D^{20} = -25.0^\circ$  ( $c = 0.02$  in  $CHCl_3$ ). {lit  $[\alpha]_D^{20} = -28.4^\circ$  ( $c = 0.32$  in  $CHCl_3$ )}.<sup>16</sup> UV  $\lambda_{CHCl_3}^{max}$  nm (log  $\epsilon$ ): 242 (3.75), 287 (3.54). MS  $m/z$  (mass/charge ratio peaks): 400 ( $M^+$ ), 265, 264, 251, 238, 219, 182, 181 (100%), 167, 135, 131, 121, 77.  $^1H$ -NMR (400 MHz,  $CDCl_3$ )  $\delta$  (chemical shift): 2.48–2.63 [(4H (protons), *m* (multiplet),  $2 \times H7'$ ,  $H8'$ , and  $H8$ )], 2.89–2.92 (2H, *m*,  $2 \times H7$ ), 3.82 [(9H, *s* (singlet),  $3 \times OMe$ )], 3.88 [(1H, *dd* (double doublet),  $J$  (coupling constant) = 9.3, 7.8 Hz,  $H9'$ )], 4.18 (1H, *dd*,  $J = 9.3$ , 7.3 Hz,  $H9'$ ), 5.93 [(1H, *d* (doublet),  $J = 1.5$  Hz,  $H10'$ )], 5.94 (1H, *d*,  $J = 1.5$  Hz,  $H10'$ ), 6.36 (2H, *s*,  $H2$  and  $H6$ ), 6.46 (1H, *d*,  $J = 1.7$  Hz,  $H2'$ ), 6.47 (1H, *dd*,  $J = 6.9$ , 1.7 Hz,  $H6'$ ), 6.69 (1H, *d*,  $J = 8.3$  Hz,  $H5'$ ). The  $^{13}C$ -NMR data are shown in Table 1.

$^1H$ -NMR and  $^{13}C$ -NMR data of compound (1) were identical with those of authentic yatein.<sup>14,15</sup>

### Podophyllotoxin

The compound (2) (podophyllotoxin) obtained from fraction 2-1 was recrystallized from chloroform and ethanol to afford white crystals (2) (14.7 mg), mp  $181^\circ$ – $183^\circ C$  (lit mp  $180.8^\circ$ – $181.8^\circ C$ ).<sup>21</sup> Compound (2) was obtained in a yield of 0.00009% of the fresh leaves. *Analysis*. Calculated:  $C_{22}H_{22}O_8$ : C, 63.75; H, 5.35. Found: C, 63.89; H, 5.36.  $[\alpha]_D^{25} = -121.3^\circ$  ( $c = 0.03$  in  $CHCl_3$ ) {lit  $[\alpha]_D^{20} = -132^\circ$  ( $c = 1.0$  in  $CHCl_3$ )}.<sup>18</sup> UV  $\lambda_{max}^{EtOH}$  nm (log  $\epsilon$ ): 219 (4.51), 291 (3.89). MS  $m/z$ : 414 ( $M^+$ ) (100%), 399, 396, 189, 181, 168, 153.  $^1H$ -NMR (400 MHz,  $CDCl_3$ )  $\delta$ : 2.17 (1H, *d*,  $J = 8.3$  Hz, OH), ca 2.7–2.9 (2H, *m*,  $H8$  and  $H8'$ ), 3.76 (6H, *s*,  $2 \times OMe$ ), 3.81 (3H, *s*, OMe), 4.08 (1H, *t* (triplet),  $J = 9.5$  Hz,  $H9'$ ), 4.59 (1H, *m*,  $H9'$ ), 4.62 (1H, *d*,  $J = 8.8$  Hz,  $H7$ ), 4.77 (1H, *t*,  $J = 8.8$  Hz,  $H7'$ ), 5.97 (1H, *d*,  $J = 1.5$  Hz,  $H10'$ ), 5.99 (1H, *d*,  $J = 1.5$  Hz,  $H10'$ ), 6.37 (2H, *s*,  $H2$  and  $H6$ ), 6.51 (1H, *s*,  $H5'$ ), 7.11 (1H, *s*,  $H2'$ ). The  $^{13}C$ -NMR data are shown in Table 1. The  $^{13}C$ -NMR spectra of compound (2) were identical with those of authentic podophyllotoxin.<sup>15</sup>

### Podophyllotoxin monoacetate

Crude compound (2) (20 mg) was acetylated with acetic anhydride (1 ml) in pyridine (1 ml), and the workup was conducted in the usual manner to afford monoacetate (3) (20.1 mg) after recrystallization with ethanol, mp  $207^\circ$ – $209^\circ C$  (lit mp  $206^\circ$ – $207^\circ C$ ).<sup>23</sup> *Analysis*. Calculated:  $C_{24}H_{24}O_9$ : C, 63.14; H, 5.30. Found: C, 63.04; H, 5.39. UV  $\lambda_{max}^{MeOH}$  nm (log  $\epsilon$ ): 209 (4.87), 278 (4.29). MS  $m/z$ : 456 ( $M^+$ ) (100%), 397, 351, 282, 229, 185, 168, 149.  $^1H$ -NMR (400 MHz,  $CDCl_3$ )  $\delta$ : 2.19 (3H, *s*, OAc), ca. 2.8–2.9 (2H, *m*,  $H8$  and  $H8'$ ), 3.76 (6H, *s*,  $2 \times OMe$ ), 3.81 (3H, *s*, OMe), 4.20 (1H, *t*,  $J = 9.7$  Hz,  $H9'$ ), 4.39 (1H, *dd*,  $J = 9.3$ , 6.8 Hz,  $H9'$ ), 4.61 (1H, *d*,  $J = 4.4$  Hz,  $H7$ ), 5.88 (1H, *d*,  $J = 8.8$  Hz,  $H7'$ ), 5.98 (1H, *d*,  $J = 1.5$  Hz,  $H10'$ ), 6.0 (1H, *d*,  $J = 1.5$  Hz,  $H10'$ ), 6.39 (2H, *s*,  $H2$  and  $H6$ ), 6.54 (1H, *s*,  $H5'$ ), 6.77 (1H, *s*,  $H2'$ ).  $^{13}C$ -NMR (100 MHz,  $CDCl_3$ )  $\delta$ : 21.1 ( $-O-CO-CH_3$ ), 38.7 ( $C8'$ ), 43.7 ( $C7$ ), 45.6 ( $C8$ ), 56.2 ( $C10$  and  $C12$ ), 60.7 ( $C11$ ), 71.3 ( $C9'$ ), 73.6 ( $C7'$ ), 101.6 ( $C10'$ ), 107.0 ( $C2'$ ), 108.1 ( $C2$  and  $C6$ ), 109.7 ( $C5'$ ), 128.3 ( $C1'$ ), 132.3 ( $C6'$ ), 134.8 ( $C1$ ), 137.2 ( $C4$ ), 147.6 ( $C3'$ ), 148.1 ( $C4'$ ), 152.6 ( $C3$  and  $C5$ ), 171.4 ( $-O-CO-CH_3$ ), 173.6 ( $C9$ ). The mixed-melting-point test of compound (3) with authentic podophyllotoxin monoacetate proved compound (3) to be identical with podophyllotoxin monoacetate.

**Acknowledgments** The authors thank Mr. Tadashi Watanabe, the chief priest of Ibuki Shinto Shrine, Uwajima City, Ehime Prefecture, for the gift of *J. chinensis* leaves.

## References

1. Tachibana S, Watanabe E, Itoh K, Oki T (1994) Formation of taxol in *Taxus cuspidata* Sieb. et Zucc. var. *nana* Rehder Callus cultures. Mokuza Gakkaishi 40:1254–1258

2. Yoshida M, Kitahata M, Itoh K, Oki T, Tachibana S (1995) Formation of isoharringtonine, an anti-tumor alkaloid in *Cephalotaxus harringtonia* Callus cultures (in Japanese). *Bull Ehime Univ For* 33:41–52
3. Tachibana S, Sumimoto M (1989) Utilization of forest resources by microbial, enzymatical, and chemical conversions. I. (in Japanese). *Mokuzai Gakkaishi* 35:42–50, 51–57
4. Tachibana S, Umemura M, Sumimoto M (1989) Utilization of forest resources by microbial, Enzymatical, and chemical conversions. II. (in Japanese). *Mokuzai Gakkaishi* 35:761–770
5. Itoh K, Tachibana S, Onogi T, Kubota M, Higashi M (1993) Production of gibberelins from (–)-kaurene with a bio-reactor by microbial conversion with *Gibbrella fujikuroi* (in Japanese). *Mokuzai Gakkaishi* 39:610–615
6. Itoh K, Tachibana S, Sumimoto M (1993) Enzymatic conversion of apigenin to 3',8-dimer type bisflavonoid (in Japanese). *Mokuzai Gakkaishi* 39:684–690
7. Itoh K, Tachibana S, Sumimoto M (1993) Enzymatic conversion of flavonoids to their methyl ethers by an extracellular crude enzyme solution from *Phanerochaete chrysosporium* (in Japanese). *Mokuzai Gakkaishi* 39:691–699
8. Uehara K (1969) *Jumoku Daizusetsu*, vol 1. Ariake Shobo, Tokyo, pp 475–488
9. Sawada T (1958) Studies on flavonoids in the leaves of coniferae and allied plants. V. (in Japanese). *Yakugaku Zasshi* 78:1023–1027
10. Fang JM, Lee CK, Cheng YS (1992) Lignans from leaves of *Juniperus chinensis*. *Phytochemistry* 31:3659–3661
11. Feliciano AS, Miguel JM, Gordaliza M, Castro A (1990) Lignans from *Juniperus sabina*. *Phytochemistry* 29:1335–1338
12. Feliciano AS, Miguel JM, Gordaliza M, Castro A (1991) Lignans from *Juniperus sabina*. *Phytochemistry* 30:3483–3485
13. McDoniel PB, Cole JR (1972) Antitumor activity of *Bursera schlechtendalii* (Burseraceae) *J Pharm Sci* 61:1992–1994
14. Kamil WM, Dewick PM (1986) Biosynthetic relationship of aryltetralin lactone lignans to dibenzylbutyrolactone lignans. *Phytochemistry* 25:2093–2102
15. Agrawal PK, Thakur RS (1986) <sup>13</sup>C NMR spectroscopy of lignan and neolignan derivatives. *Magn Reson Chem* 23:389–418
16. Erdtman H, Harmatha J (1979) Phenolic and terpenoid heartwood constituents of *Libocedrus yateensis*. *Phytochemistry* 18:1495–1500
17. Jackson DE, Dewick PM (1984) Aryltetralin lignans from *Podophyllum hexandrum* and *Podophyllum peltatum*. 25:1147–1152
18. Hartwell JL, Schrecker AW (1951) Components of podophyllin. V. *J Am Chem Soc* 73:2909–2916
19. Hartwell JL, Detty WE (1950) Components of podophyllin. III. 72:246–253
20. Hartwell JL (1976) Types of anticancer agents isolated from plants. *Cancer Treat Rep* 60:1031–1067
21. Jackson DE, Dewick PM (1985) Tumour-inhibitory aryltetralin lignans from *Podophyllum pleianthum*. *Phytochemistry* 24:2407–2409
22. Miyata M, Muranaka T, Itoh K, Oki T, Tachibana S (1996) Formation of podophyllotoxin in *Juniperus chinensis* L. callus cultures (in Japanese). In: Abstracts of the 46th annual meeting of the Japan Wood Research Society, Kumamoto, p 497
23. Nadkarni MV, Hartwell JL, Maury PB, Leiter J (1953) Components of podophyllin. IX. *J Am Chem Soc* 75:1308–1312